

AD-A055 033

TEXAS UNIV AT AUSTIN CENTER FOR CYBERNETIC STUDIES
GRADIENT STATES FOR SOME DUALITIES WITH THE CHARNES-COOPER EXTR--ETC(U)
JUL 77 A CHARNES, L SEIFORD
CCS-295

F/G 12/1

N00014-75-C-0569


NL

UNCLASSIFIED

1 OF 1
AD
A055 033



END
DATE
FILMED
7-78
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 055033


FOR FURTHER TRAN ¹¹⁻¹¹

12



CENTER FOR CYBERNETIC STUDIES

The University of Texas
Austin, Texas 78712



This document has been approved
for public release and sale; its
distribution is unlimited.



78 06 06 056


12

Research Report CCS 295

GRADIENT STATES FOR SOME DUALITIES
WITH THE C^2 EXTREMAL PRINCIPLE

by

A. Charnes*
L. Seiford**



September 1977

*The University of Texas, Austin, Texas.
**York University, Downsview, Ontario, Canada.

This research was partly supported by Project NR047-021, CNR Contract N00014-75-C-0569 with the Center for Cybernetic Studies, The University of Texas. Reproduction in whole or in part is permitted for any purpose of the United States Government.

CENTER FOR CYBERNETIC STUDIES

A. Charnes, Director
Business-Economics Building, 203E
The University of Texas
Austin, Texas 78712
(512) 471-1821

78 06 06 056

ABSTRACT

Gradient characterizations of some convex function infima are derived which apply to extension of the Charnes-Cooper duality state characterizations to more general classes of convex programming problems via the Charnes-Cooper extremal principle for optimization dualities.

| | |
|---------------------------------|---|
| ACCESSION for | |
| NTIS | White Section <input checked="" type="checkbox"/> |
| DDC | Buff Section <input type="checkbox"/> |
| UNANNOUNCED | <input type="checkbox"/> |
| JUSTIFICATION | |
| BY | |
| DISTRIBUTION/AVAILABILITY CODES | |
| Dist. | SPECIAL |
| A | |

78 06 06 056

The C^2 extremal principle for dualities which was originally presented in Charnes, Cooper and Seiford [1] is an approach to deriving dual optimization problems with proper duality inequality which simplifies and generalizes the Fenchel-Rockafellar scheme [2, 3]. The derivation is accomplished in two stages. The first is the achievement of the duality inequality. The second is the decoupling of the primal and dual variables.

The C^2 Extremal Principle

Let $K(\delta, x)$ be a real valued function which is concave in δ for

$$(\delta, x) \in \Delta \Theta X \subseteq R^m \Theta R^s$$

and for which

$$g(\delta) \equiv \inf_{x \in X} K(\delta, x)$$

exists for each $\delta \in \Delta$. Let T be a map from the convex set $Z \subseteq R^n$ into X . If $K(\delta, T(z)) = f(z)$, a convex function for $z \in Z$, $\delta \in \Gamma$, then $\Gamma \Theta T[Z]$ is the decoupling set for (δ, x) . If we further require $\Delta \cap \Gamma$ to be a convex set, the problems

$$\sup g(\delta), \quad \delta \in \Delta \cap \Gamma \tag{1}$$

and

$$\inf f(z), \quad z \in Z \tag{2}$$

are dual convex programming problems.

As an example in the use of the use of the C^2 extremal principle, we derive the dual problems for linear programming. Let

$$K(y, x) = y^T x$$

for

$$y \in \Delta = \{y : y \geq 0\}$$

and

$$x \in X = \{x : x \geq b\}.$$

Then

$$g(y) = \inf_x K(y, x) = y^T b.$$

If we let $x = Az$, then

$$y^T b \leq y^T Az \quad \forall y \in \Delta, \quad \forall z \in Z = \{z : Az \geq b\}.$$

Defining

$$\Gamma = \{y : y^T A = c^T\}$$

we have

$$\sup_{y \in \Delta \cap \Gamma} y^T b \leq \inf_{z \in Z} c^T z$$

or equivalently

$$\sup b^T y \leq \inf c^T z$$

subject to

$$A^T y = c$$

$$y \geq 0$$

subject to

$$Az \geq b$$

(3)

It is well known that a duality gap cannot occur in linear programming. In the general case the existence or non-existence of duality gaps is dependent on the choices of $\Delta \cap \Gamma$ and Z .

Gradient Characterization of Some Convex Function Infima

The extension of our characterization to duality states for more general cases of dual convex problems of the form given in Charnes, Cooper and Seiford [1] depends on developing properties characterizing the existence or non-existence of infima for special classes of convex functions. In the following theorem we adduce some such properties.

Theorem 1: Let $f: X \rightarrow \mathbb{R}$ be convex and differentiable on an open convex set $X \subseteq \mathbb{R}^m$. For the linear function $A: Z \rightarrow X$ with Z convex, consider

$$C(z) = f(Az) - b^T z.$$

If we define

$$\Gamma = \{\delta: \delta^T A = b^T\}$$

$$\Delta_x = \nabla f[x]$$

$$\Delta_z = \nabla f[A(Z)]$$

then

- (1) a) $C(z)$ is bounded below implies

$$\bar{\Delta}_z \cap \Gamma \neq \emptyset$$

- b) $\Delta_x \cap \Gamma \neq \emptyset$ implies $C(z)$ is bounded below

(ii) a) $C(z)$ has an infimum implies

$$\bar{\Delta}_z \cap \Gamma \neq \emptyset, \Delta_z \cap \Gamma = \emptyset$$

b) $\Delta_x \cap \Gamma \neq \emptyset, \Delta_z \cap \Gamma = \emptyset$ implies $C(z)$ has an infimum

(iii) $C(z)$ has a minimum if and only if $\Delta_z \cap \Gamma \neq \emptyset$.

Proof: (i) a) Suppose $\bar{\Delta}_z \cap \Gamma = \emptyset$. Then

$$\nabla C(z) = \nabla f(Az)^T A - b^T$$

is bounded away from zero, i.e.,

$$\|\nabla C(z)\| \geq \epsilon > 0.$$

Consider the differential equation system.

$$\dot{z}(t) = \frac{-\nabla C(z(t))}{\|\nabla C(z(t))\|}.$$

The function $-\nabla C(\cdot)/\|\nabla C(\cdot)\|$ is continuous [2] and bounded, since ∇C is the gradient of a convex function. Hence there exists a solution, $z(t)$. For $F(t) \equiv C(z(t))$

$$\begin{aligned} F'(t) &= \nabla C(z(t)) \cdot \dot{z}(t) \\ &= \nabla C(z(t)) \cdot \left(\frac{-\nabla C(z(t))}{\|\nabla C(z(t))\|} \right) \\ &= -\|\nabla C(z(t))\| \leq -\epsilon < 0. \end{aligned}$$

Thus as $t \rightarrow +\infty$

$$F(t) = C(z(t)) \rightarrow -\infty$$

and C is unbounded below.

(i) b) If $\Delta_x \cap \Gamma \neq \emptyset$, let $\bar{\delta} \in \Delta_x \cap \Gamma$.

$$\text{Then } C(z) = f(Az) - b^T z = f(Az) - \bar{\delta}^T Az.$$

$$\text{Hence } \inf_z C(z) \geq \inf_x f(x) - \bar{\delta}^T x \geq f(x_0) - \bar{\delta}^T x_0$$

by the differentiable convexity of f , where x_0 satisfies $\nabla f(x_0) = \bar{\delta}$.

(iii) Suppose $C(z)$ attains its minimum at z_0 . Then

$$\nabla C(z_0)^T = \nabla f(Az_0)^T A - b^T = 0.$$

Setting $\delta = \nabla f(Az_0)$ we have

$$\delta \in \Delta_z \cap \Gamma.$$

Conversely, if $\delta_0 \in \Delta_z \cap \Gamma$, then $\delta_0 \in \Gamma$

$$\text{so } C(z) = f(Az) - \delta_0^T Az$$

$$\begin{aligned} \text{and } \nabla C(z)^T &= \nabla f(Az)^T \cdot A - \delta_0^T A \\ &= (\nabla f(Az))^T - \delta_0^T A. \end{aligned}$$

Since $\delta_0 \in \Delta_z$, $\exists z_0$ and that $\nabla f(Az_0) = \delta_0$.

Hence $\nabla C(z_0) = 0$ and $C(z)$ attains its minimum at z_0 .

(ii) a) and b) now follow by exhaustion.

Corollary 1 $\Gamma = \emptyset \Rightarrow C(z)$ is unbounded below.

Proof: Consider the dual linear programming problems

$$\begin{array}{ll} \text{(I)} & \text{(II)} \\ \max & b^T z & \min & \delta^T 0 \\ \text{s.t.} & Az = 0 & \text{s.t.} & \delta^T A = b^T \end{array}$$

If $\Gamma = \emptyset$, then II is infeasible. Since $z = 0$ satisfies (I), there exists a sequence z_n such that

$$Az_n = 0 \quad (\forall n) \text{ and } b^T z_n \uparrow +\infty.$$

$$\begin{aligned} \text{Thus } C(z_n) &= f(Az_n) - b^T z_n \\ &= f(0) - b^T z_n \rightarrow -\infty. \end{aligned}$$

That the characterization given by Theorem 1 is a best possible is shown by the following examples.

Example 1: To show that (in i, b) we need $\Delta_x \cap \Gamma \neq \emptyset$ (rather than $\bar{\Delta}_x \cap \Gamma \neq \emptyset$) to insure $C(z)$ is bounded below, consider

$$f(x) = \begin{cases} -\ln(-x) & \text{if } x \leq -1 \\ x + 1 & \text{if } x > -1 \end{cases}$$

Then $b = 0 \in \bar{\Delta}_x$ but $\lim_{x \rightarrow -\infty} f(x) - 0 \cdot x = -\infty$.

Example 2: To show (in i, a) that $C(z)$ bounded below only guarantees $\bar{\Delta}_z \cap \Gamma \neq \emptyset$ and not $\Delta_z \cap \Gamma \neq \emptyset$, consider

$$f(z) = e^z.$$

Then $f(z) = e^z - 0 \cdot z > 0 \quad (\forall z)$

but $0 \notin \Delta_z, 0 \in \bar{\Delta}_z$.

Conclusion

In other work now in progress we employ these results to obtain duality state characterizations of dual convex programs derived from the C^2 principle. We also make applications to two-person zero-sum games whose payoff function is of the form $K(\delta, x) = f(x) - \delta^T x + g(\delta)$ where $f(x)$ is convex and $g(\delta)$ is concave. Such

games have arisen in contexts where the x -player corresponds to a government agency and the δ -player is the totality of enterprise groups whose activities are being regulated.

1. A. Charnes, W. W. Cooper and L. Hellwig, "Extremal Principles and Optimization Problems for Linear-Quadratic-Linear Problems," Center for Cybernetic Studies, Research Report CCS 161, The University of Texas, Austin, Texas, April 1976. To appear in *Mathematical Operations Research and Statistics*, Series Optimization, Issue 1, vol. 9 (1978).

2. W. Fenchel, "Convex Cones, Sets, and Functions," Lecture Notes, Princeton University, Department of Mathematics, September 1953.

3. E. T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, New Jersey, 1970.

REFERENCES

1. A. Charnes, W. W. Cooper and L. Seiford, "Extremal Principles and Optimization Dualities for Khinchin-Kullback-Leibler Estimation," Center for Cybernetic Studies, Research Report CCS 261, The University of Texas, Austin, Texas, April 1976. To appear in *Zeitschrift Mathematische Operationsforschung und Statistik, Series Optimization*, issue 1, vol. 9 (1978).
2. W. Fenchel, "Convex Cones, Sets, and Functions," Lecture Notes, Princeton University, Department of Mathematics, September 1953.
3. R. T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, New Jersey, 1970.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

| | | | |
|---|--|---|----------------------|
| 1. ORIGINATING ACTIVITY (Corporate author) Center for Cybernetic Studies The University of Texas | | 2a. REPORT SECURITY CLASSIFICATION Unclassified | |
| | | 2b. GROUP | |
| 3. REPORT TITLE ⑥ Gradient States for Some Dualities with the ② Extremal Principle. | | | |
| 4. DESCRIPTIVE NOTES (Type of report and, inclusive dates) ⑩ Charnes - Cooper | | | |
| 5. AUTHOR(S) (First name, middle initial, last name) A./Charnes L./Seiford | | | |
| 6. REPORT DATE Jul 77 ⑪ | | 7a. TOTAL NO. OF PAGES ⑫ 13 P. | 7b. NO. OF REFS 3 |
| 8a. CONTRACT OR GRANT NO. N00014-75-C-0569 ⑮ | | 9a. ORIGINATOR'S REPORT NUMBER(S) ⑨ Center for Cybernetic Studies Research Report CCS 295 | |
| b. PROJECT NO. NR047-021 | | 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) | |
| 10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited. | | | |
| 11. SUPPLEMENTARY NOTES | | 12. SPONSORING MILITARY ACTIVITY Office of Naval Research (Code 434) Washington, D.C. | |
| 13. ABSTRACT Gradient characterizations of some convex function infima are derived which apply to extension of the Charnes-Cooper duality state characterizations to more general classes of convex programming problems via the Charnes-Cooper extremal principle for optimization dualities. | | | |

Unclassified

Security Classification

A-31408
hc

Unclassified

Security Classification

| 14 KEY WORDS | LINK A | | LINK B | | LINK C | |
|-----------------------------------|--------|----|--------|----|--------|----|
| | ROLE | WT | ROLE | WT | ROLE | WT |
| Gradient characterizations | | | | | | |
| Convex function infima | | | | | | |
| Charnes-Cooper extremal principle | | | | | | |
| Dual convex programs | | | | | | |

DD FORM 1473 (BACK)
1 NOV 65
5/N 0102-014-6800

Unclassified

Security Classification

A-1403